arXiv:astro-ph/0011388v1 21 Nov 2000

# METAL ABUNDANCE OF THE ECLIPSING BINARY YZ CAS

*Discrepancy between atmospheric determinations for the Am component, YZ Cas A*

E. Lastennet
*Astronomy Unit, QMW, Mile End Road, London E1 4NS, UK*

D. Valls-Gabaud
*Lab. d'Astrophysique, UMR CNRS 5572, OMP, 14 av. Belin, F-31400 Toulouse, France*

C. Jordi
*Dep. d'Astronomia i Meteorologia, Av. Diagonal 647, E-08028 Barcelona, Spain*

**Abstract**

We review current and new estimates of the eclipsing binary YZ Cassiopeiae metallicity (Z). Since the individual components cover a quite large range of mass (1.35-2.31$M_\odot$), YZ Cas is potentially one of the best stellar laboratories to understand the structure and evolution of 1 to 2 $M_\odot$ stars. The derivation of Z from IUE spectra, as well as from photometric indices, provides the chemical composition of the atmosphere ($Z_{\rm atmospheric}$), while the fit of evolutionary tracks provides the initial chemical composition ($Z_{\rm initial}$). While a disagreement is expected between $Z_{\rm atmospheric}$ and $Z_{\rm initial}$ because the primary component is an Am star (one expects $Z_{\rm atmospheric}$ to be larger), we find some unexpected discrepancy between atmospheric determinations of Z for this star.

## 1. $Z_{\rm ATMOSPHERIC}$ FROM IUE SPECTRA

From IUE spectra of the YZ Cas primary, [LM83] found that the metals add up to a Z value of over 0.03 (Z=0.036±0.005). We try to assess to what extent this high Z-value has to be revised. It is well known that the abundance of iron is enhanced in Am stars with respect to normal A-F stars. [LM83] found for YZ Cas A that Fe is ∼6 times

overabundant, which implies that Fe contributes to ∼43% of Z. In order to update the [LM83] study, we applied two corrections to this value: 1) They assumed an iron solar abundance $A_{Fe}$=7.60. More detailed studies give $A_{Fe}$ = 7.50±0.05 ([GS99]) implying that Fe contributes to ∼57% of Z. 2) Their analysis assumed $T_{eff}$=10,300K. Recent determinations give 9100±300K ([R00]). This would reduce the [LM83]'s iron abundance by about 0.15-0.20 dex, and then imply that Fe contributes to ∼33% of Z. Both corrections are opposite and imply altogether that Fe contributes to ∼40% of Z. Hence, our attempt to update some of the [LM83] assumptions does not change their high-Z conclusion.

## 2. $Z_{ATMOSPHERIC}$ FROM PHOTOMETRIC CALIBRATIONS

Since *uvbyβ* Strömgren colours are available for both components of YZ Cas ([L81], [J97]), it is possible to derive [Fe/H] from existing calibrations. The [J97] photometric indices do not suggest a metallic behaviour of the Am star atmosphere: [Fe/H] is −0.02±0.22 ([J00]), i.e $Z=0.017^{+0.012}_{-0.007}$ (assuming $Z_{\odot}$=0.018). The upper limit is only marginally compatible with the Z derived from the IUE spectra. Assuming the individual photometric indices from [L81], the derived atmospheric [Fe/H] is 0.12±0.05, i.e. Z=0.024±0.003, again only marginally compatible with the IUE determination. Using Strömgren photometry from [J97] and accurate gravities, [L99] derived simultaneous $T_{eff}$-[Fe/H] estimates from the BaSeL photometric calibrations ([L98]). They obtain [Fe/H] strictly lower than 0.1 for YZ Cas A, i.e Z<0.022. This result assumes no reddening which is justified by experiments we performed with the BaSeL models (see [L99] for details on the method) for E(b−y) ranging from −0.034 (*negative* value of [L81]) to 0.09 (maximum value quoted by [LM83]). In summary, even if the results of [J00] and [L99] are not completely independent because based on the same colour indices, they suggest that the atmospheric metallicity of YZ Cas A is solar or sub-solar, and exclude Z-values greater than 0.03.

## References

De Landtsheer A.C., Mulder P.S. 1983, A&A 127, 297 [LM83]
Grevesse N., Sauval A.J. 1999, A&A 347, 348 [GS99]
Jordi C., Ribas I., Torra J., Giménez A. 1997, A&A 326, 1044 [J97]
Jordi C., 2000, private communication [J00]
Lacy C.H. 1981, ApJ 251, 591 [L81]
Lastennet E., Lejeune Th., Westera P., Buser R. 1999, A&A 341, 857 [L99]
Lejeune Th., Cuisinier F., Buser R. 1998, A&AS 130, 65 [L98]
Ribas I., Jordi C., Torra J., Giménez A. 2000, MNRAS 313, 99 [R00]